

BMJ Best Practice

Hypercalcaemia of malignancy

Straight to the point of care



Last updated: Apr 08, 2021

Table of Contents

Overview	3
Summary	3
Definition	3
Theory	4
Epidemiology	4
Aetiology	4
Pathophysiology	4
Classification	6
Case history	7
Diagnosis	8
Approach	8
History and exam	12
Risk factors	13
Investigations	16
Differentials	18
Criteria	19
Management	20
Approach	20
Treatment algorithm overview	22
Treatment algorithm	23
Patient discussions	29
Follow up	30
Monitoring	30
Complications	30
Prognosis	30
Guidelines	32
Diagnostic guidelines	32
Treatment guidelines	32
References	34
Images	37
Disclaimer	39

Summary

Hypercalcaemia of malignancy can result from: humoral hypercalcaemia of malignancy (characterised by tumour secretion of parathyroid hormone-related peptide [PTHrP]); local osteolytic hypercalcaemia (characterised by local release of factors, including PTHrP, by bony metastases that promote osteoclast differentiation and function); calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia (characterised by autonomous production of calcitriol [(1,25-dihydroxyvitamin D)] by lymphoma cells); and ectopic hyperparathyroidism (characterised by tumour production of parathyroid hormone), which is very rare.

Pharmacological therapy can transiently improve hypercalcaemia. However, sustained maintenance of normocalcaemia requires eradication of underlying malignancy.

Intravenous bisphosphonates are the most studied and efficacious agents to treat humoral hypercalcaemia of malignancy and local osteolytic hypercalcaemia.

Glucocorticoids may be efficacious for treating calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia.

Definition

Hypercalcaemia occurs in 20% to 30% of patients with cancer.[1] [2] Cancer represents the most common aetiology of hypercalcaemia in the inpatient setting.[3] It can result from: humoral hypercalcaemia of malignancy (characterised by tumour secretion of parathyroid hormone-related peptide [PTHrP]); local osteolytic hypercalcaemia (characterised by local release of factors, including PTHrP, by bony metastases that promote osteoclast differentiation and function); calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia (characterised by autonomous production of calcitriol [(1,25-dihydroxyvitamin D)] by lymphoma cells); and ectopic hyperparathyroidism (characterised by tumour production of parathyroid hormone), which is very rare.[1] [4]

Epidemiology

Hypercalcaemia occurs in 20% to 30% of patients with cancer.^{[1] [2]} Cancer represents the most common aetiology of hypercalcaemia in the inpatient setting.^[3] Humoral hypercalcaemia of malignancy (parathyroid hormone-related peptide [PTHrP]-mediated) accounts for 80% of cases, with typical tumours including squamous cell cancer, renal cancer, ovarian cancer, endometrial cancer, breast cancer, and human T-lymphotrophic virus-associated lymphoma.^{[1] [2] [4] [5]} Local osteolytic hypercalcaemia (local production of factors, including PTHrP, by bony metastases that promote osteoclast differentiation and activity) accounts for 20% of cases, with typical tumours including breast cancer, multiple myeloma, and lymphoma.^{[1] [4] [5]} Hypercalcaemia mediated by increased calcitriol (1,25-dihydroxyvitamin D) synthesis or ectopic parathyroid hormone secretion accounts for <1% of cases.^{[1] [5]}

Aetiology

Humoral hypercalcaemia of malignancy is most commonly associated with:^{[1] [2] [4] [5]}

- Renal cancer
- Ovarian cancer
- Breast cancer
- Endometrial cancer
- Human T-lymphotrophic virus-associated lymphoma
- Squamous cell carcinoma.

Local osteolytic hypercalcaemia is most commonly associated with:^{[1] [4] [5]}

- Breast cancer
- Multiple myeloma


Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia is associated with lymphomas of all types, but is also seen with granulomatous disease (such as active sarcoidosis) or tuberculosis.^{[1] [5]}

Ectopic parathyroid hormone secretion by tumours is rare, but has been reported.^{[1] [4] [7]}

Pathophysiology


Humoral hypercalcaemia of malignancy: tumour secretion of parathyroid hormone-related peptide (PTHrP) leads to activation of osteoclastic bone resorption and suppression of osteoblastic bone formation. This results in skeletal release of calcium and subsequent hypercalcaemia. PTHrP also acts at the level of the kidney to reduce calcium clearance, as well as to reduce the renal phosphorus threshold, leading to hyperphosphaturia and hypophosphataemia.^{[1] [2] [4]}

Local osteolytic hypercalcaemia: in the setting of malignancy with widespread skeletal involvement, local release of cytokines (e.g. IL-1 and IL-6), chemokines, and PTHrP leads to increased osteoclastic bone resorption, and to increased release of calcium into the serum that overwhelms the kidney's capacity to clear calcium.^[4]



CT chest showing compression fracture of multiple vertebral bodies in a child presenting with acute lymphoblastic leukaemia. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level

Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758



Whole body planar images suggestive of skeletal infiltration in a child with acute lymphoblastic leukaemia showing areas of abnormal increased uptake. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level
 Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. *BMJ Case Reports* 2013; doi:10.1136/bcr-2013-008758

Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia: overexpression of 1-alpha hydroxylase (the enzyme that converts precursor 25-hydroxyvitamin D [25OHD] to calcitriol [(1,25-dihydroxyvitamin D)]) by malignant or adjacent normal cells leads to autonomous production of bioactive vitamin D.[1] [4] Calcitriol (1,25-dihydroxyvitamin D) in turn increases intestinal calcium absorption and subsequent hypercalcaemia. In addition, calcitriol (1,25-dihydroxyvitamin D) may stimulate osteoclast-mediated bone resorption.[4]

Ectopic secretion of authentic parathyroid hormone with subsequent hypercalcaemia has been rarely reported.[1] [4] [7]

Classification

Severity

While there are no formal classification schemes for defining severity of hypercalcaemia, some consider mild hypercalcaemia to be a total adjusted serum calcium of 2.6 to 3.0 mmol/L (10.5 to 12.0 mg/dL), moderate hypercalcaemia a level of 3.0 to 3.4 mmol/L (12.0 to 13.7 mg/dL), and severe hypercalcaemia a level greater than 3.4 mmol/L (13.7 mg/dL).[1] [5] [6]

Causes of hypercalcaemia of malignancy

Humoral hypercalcaemia of malignancy is most commonly associated with:[1] [4] [5]

- Renal cancer
- Ovarian cancer
- Breast cancer
- Endometrial cancer
- Human T-lymphotrophic virus-associated lymphoma
- Squamous cell carcinoma.

Local osteolytic hypercalcaemia is most commonly associated with:[1] [4] [5]

- Breast cancer
- Multiple myeloma

Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia is associated with lymphomas of all types, but is also seen with granulomatous disease (such as active sarcoidosis) or tuberculosis.[1] [5]

Ectopic parathyroid hormone secretion by tumours is rare, but has been reported.[1] [4] [7]

Case history

Case history #1

A 63-year-old woman is brought in by her family for progressive fatigue and confusion. Past medical history is notable for ovarian cancer. Physical examination reveals dry mucous membranes. Admission labs are significant for an elevated adjusted serum calcium of 3.2 mmol/L (12.8 mg/dL), a low-normal albumin level, a low-normal phosphorus level, and elevated alkaline phosphatase. Hypercalcaemia work-up reveals a suppressed parathyroid hormone, an elevated parathyroid hormone-related peptide (PTHrP), and a low-normal calcitriol (1,25-dihydroxyvitamin D) level.

Other presentations

Hypercalcaemia of malignancy can occur with both solid and liquid tumours. Presenting symptoms are similar to those seen in hypercalcaemia of any aetiology. Symptoms include neuropsychiatric changes (mood disturbance, fatigue, confusion, stupor, and coma), gastrointestinal disturbance (loss of appetite, nausea, constipation), muscle weakness, acute kidney injury, polyuria, polydipsia, and bone pain.[1] [4] [6] Mild hypercalcaemia may be asymptomatic. Severe hypercalcaemia may be associated with hypercalcaemic crisis, with complications such as acute pancreatitis, acute kidney injury, and coma.[5]

Approach

Diagnosis of malignancy-associated hypercalcaemia is based on a history of malignancy, signs/symptoms of hypercalcaemia, presence of hypercalcaemia, and biochemical work-up to determine the underlying aetiology of hypercalcaemia.

History and physical examination

Symptoms of hypercalcaemia include neuropsychiatric changes (e.g., mood disturbance, fatigue, confusion, stupor, and coma), gastrointestinal disturbance (loss of appetite, nausea, constipation), muscle weakness, polyuria, polydipsia, and bone pain.[1] [2] [4] [5] [6] Signs of dehydration associated with hypercalcaemia include poor skin turgor and dry mucous membranes. However, hypercalcaemia may not be associated with any specific physical examination findings.

History to evaluate potential causes of hypercalcaemia unrelated to malignancy should be elicited.[1] [4] These include hyperthyroidism, pheochromocytoma, adrenal insufficiency, and granulomatous disease. A thorough medication history to evaluate for use of medications that may cause or worsen hypercalcaemia (thiazide diuretics, lithium, calcium, over-the-counter antacids, and large doses of vitamin D) should be undertaken.[1] [4] [5]

Severity of symptoms may correlate with duration and rapidity of onset of hypercalcaemia.[5] It is important to note that malignancy-associated hypercalcaemia is less commonly associated with life-threatening decompensation, but a hypercalcaemic crisis (characterised by oliguria, anuria, somnolence, coma) can occur.

An ECG should be performed to assess for a shortened QT interval and dysrhythmias.

Laboratory evaluation

Biochemical evidence of hypercalcaemia should be demonstrated by measuring total serum calcium and albumin in order to calculate the adjusted serum calcium level. Some sources prefer measurement of serum ionised calcium, if available. Serum ionised calcium should be measured if the albumin level is altered, and if calcium-binding immunoglobulins may be present (e.g., in multiple myeloma).[1]

Parathyroid hormone (PTH)-dependent hypercalcaemia should be ruled out by measuring a serum intact PTH level.[4] Conditions associated with elevated PTH include primary and tertiary hyperparathyroidism, familial hypocalciuric hypercalcaemia, and ectopic hyperparathyroidism.


A complete metabolic panel should be ordered to check for hypercalcaemia-induced acute kidney injury. A high bicarbonate level may indicate that malignancy or exogenous calcium excess may be a more likely cause of hypercalcaemia than primary hyperparathyroidism.

The aetiology of non-PTH-dependent hypercalcaemia should be determined. Conditions associated with hypercalcaemia and an appropriately suppressed PTH include malignancy-associated hypercalcaemia, hyperthyroidism, pheochromocytoma, and adrenal insufficiency.


All laboratory tests should be obtained before treatment is started, as improvement of hypercalcaemia after treatment is initiated may cause difficulty in interpreting test results. The following tests are also recommended:

- Screening for hyperthyroidism, pheochromocytoma, and adrenal insufficiency, as clinically indicated.
- Serum parathyroid hormone-related peptide (PTHrP). An elevated PTHrP associated with a suppressed PTH and low to low-normal calcitriol (1,25-dihydroxyvitamin D) level represents likely humoral hypercalcaemia of malignancy. PTHrP-mediated hypercalcaemia typically occurs with non-metastatic tumours that may be apparent on examination and initial imaging. Exceptions include small neuro-endocrine tumours.[1]
- Serum phosphorus. In humoral hypercalcaemia of malignancy, PTHrP also acts at the level of the kidney to reduce calcium clearance, as well as to reduce the renal phosphorus threshold, leading to hyperphosphaturia and hypophosphataemia.[1] [4]
- Calcitriol (serum 1,25-dihydroxyvitamin D) level, if lymphoma and/or granulomatous disease are suspected. Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia occurs with lymphomas of all kinds and granulomatous disorders.
- Serum 25-hydroxyvitamin D, if intravenous bisphosphonate or subcutaneous denosumab is being considered. Vitamin D deficiency should be corrected prior to administration of bisphosphonates or denosumab to avoid the risk of hypocalcaemia and, possibly, osteonecrosis of the jaw.[11] [12] [13] [14]

If there is a history of malignancy complicated by skeletal involvement, PTHrP and calcitriol (1,25-dihydroxyvitamin D) are found to be normal, and other aetiologies of hypercalcaemia have been eliminated, local osteolytic hypercalcaemia is the most likely diagnosis.



CT chest showing compression fracture of multiple vertebral bodies in a child presenting with acute lymphoblastic leukaemia. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level
Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758



Whole body planar images suggestive of skeletal infiltration in a child with acute lymphoblastic leukaemia showing areas of abnormal increased uptake. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level
Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758

Radiology evaluation

In order to assess for underlying causes of hypercalcaemia, the following tests should be considered:

- Skeletal survey in patients in whom multiple myeloma, bone metastases, or leukaemia are suspected.
- Chest x-ray to assess for lung cancer, sarcoidosis, or tuberculosis.

Further imaging should be ordered, according to the suspected cancer or condition and the anatomical region involved.

Other tests

An ECG should be performed to look for shortened QT interval or other conduction abnormalities.[2]

History and exam

Key diagnostic factors

presence of risk factors (common)

- Key risk factors include non-metastatic malignancy (humoral hypercalcaemia), metastatic skeletal involvement (local osteolytic hypercalcaemia), and lymphoma (calcitriol [(1,25-dihydroxyvitamin D)]-mediated hypercalcaemia).

history of malignancy (common)

- Hypercalcaemia occurs in 20% to 30% of patients with cancer.[1] [2] Cancer represents the most common aetiology of hypercalcaemia in the inpatient setting.[3] It can result from: humoral hypercalcaemia of malignancy (characterised by tumour secretion of parathyroid hormone-related peptide [PTHrP]); local osteolytic hypercalcaemia (characterised by local release of factors, including PTHrP, by bony metastases that promote osteoclast differentiation and function); calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia (characterised by autonomous production of calcitriol [(1,25-dihydroxyvitamin D)] by lymphoma cells); and ectopic hyperparathyroidism (characterised by tumour production of parathyroid hormone), which is very rare.[1] [4]

Other diagnostic factors

normal physical exam (common)

- Hypercalcaemia may not be associated with any specific physical examination findings.

poor skin turgor and/or dry mucous membranes (common)

- Signs of dehydration may be apparent on physical examination.

confusion (common)

- Hypercalcaemia may be associated with neuropsychiatric symptoms.

fatigue (common)

- Hypercalcaemia may be associated with neuropsychiatric symptoms.

constipation (common)

- Hypercalcaemia is associated with gastrointestinal symptoms.

loss of appetite (common)

- Hypercalcaemia is associated with gastrointestinal symptoms.

nausea (common)

- Hypercalcaemia is associated with gastrointestinal symptoms.

polyuria (common)

- Hypercalcaemia is associated with increased urinary excretion and dehydration by inducing nephrogenic diabetes insipidus.

polydipsia (common)

- Hypercalcaemia is associated with increased urinary excretion and dehydration.

bone pain (common)

- Hypercalcaemia is associated with bone pain.
- Bone pain may be a feature of metastatic skeletal involvement.

use of hypercalcaemia-inducing medication (common)

- Medications that may cause or worsen hypercalcaemia include thiazide diuretics, lithium, calcium, over-the-counter antacids, and large doses of vitamin D.[1] [4] [5]

stupor (uncommon)

- Hypercalcaemia may be associated with neuropsychiatric symptoms. Stupor is a non-specific symptom that may result from hypercalcaemia, but has numerous toxic/metabolic and neurological aetiologies that need to be considered.

coma (uncommon)

- Hypercalcaemia may be associated with neuropsychiatric symptoms. Coma is a non-specific symptom that may result from hypercalcaemia, but has numerous toxic/metabolic and neurological aetiologies that need to be considered.

Risk factors


Strong

non-metastatic malignancy


- Humoral hypercalcaemia of malignancy results from tumour secretion of parathyroid hormone-related peptide (PTHrP), and typically occurs with non-metastatic disease.[8] Types of cancers include renal cancer, ovarian cancer, breast cancer, endometrial cancer, and squamous cell carcinoma.[8] [9] [10]

metastatic skeletal involvement

- Local osteolytic hypercalcaemia occurs with disease complicated by widespread skeletal involvement.[8] Types of cancers include breast cancer and multiple myeloma.[8] [9] [10]



CT chest showing compression fracture of multiple vertebral bodies in a child presenting with acute lymphoblastic leukaemia. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level
Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758



Whole body planar images suggestive of skeletal infiltration in a child with acute lymphoblastic leukaemia showing areas of abnormal increased uptake. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level
Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758

lymphoma

- Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia occurs with lymphomas of all types, including human T-lymphotrophic virus-associated lymphoma.[8] [10]

Investigations

1st test to order

Test	Result
<p>total serum calcium</p> <ul style="list-style-type: none"> Hypercalcaemia is defined by an elevated serum calcium level. Total serum calcium measurements are influenced by serum albumin levels, and by calcium-binding immunoglobulins as in multiple myeloma.[1] Serum calcium measurements should ideally be obtained from specimens collected without the use of a tourniquet/cuff. May be included in comprehensive metabolic panel. 	elevated
<p>serum ionised calcium</p> <ul style="list-style-type: none"> Some sources prefer measurement of serum ionised calcium to total serum calcium plus serum albumin, if available. Measurement of serum ionised calcium should be considered when factors that affect the accuracy of total serum calcium are present, such as abnormal serum albumin levels, and calcium-binding immunoglobulins as in multiple myeloma.[1] Serum calcium measurements should ideally be obtained from specimens collected without the use of a tourniquet/cuff. 	elevated
<p>serum albumin</p> <ul style="list-style-type: none"> For SI units (calcium in mmol/L; albumin in g/L), the formula for adjusted serum calcium is $(0.02 \times [\text{normal albumin} - \text{patient's albumin}]) + \text{serum calcium}$. Calculation of albumin-adjusted total calcium is not required when ionised calcium is measured. The formula for adjusted calcium may not be reliable in all circumstances (such as in critically ill patients). May be included in comprehensive metabolic panel. 	variable
<p>comprehensive metabolic panel</p> <ul style="list-style-type: none"> May show elevated urea and creatinine in acute kidney injury; elevated bicarbonate level may indicate malignancy or exogenous calcium excess as a cause for hypercalcaemia. 	may show elevated urea or bicarbonate
<p>resting ECG</p> <ul style="list-style-type: none"> A resting ECG should be done to check the QT interval or other conduction abnormalities. 	may show shortened QT interval and dysrhythmias
<p>serum intact parathyroid hormone</p> <ul style="list-style-type: none"> Should be ordered for initial biochemical assessment of hypercalcaemia to distinguish between parathyroid hormone (PTH)-mediated hypercalcaemia and non-PTH-mediated hypercalcaemia. Malignancy-associated hypercalcaemia generally leads to appropriate suppression of PTH secretion by the parathyroid gland. However, co-existing malignancy and primary hyperparathyroidism should be excluded. Rare scenarios in which PTH is inappropriately normal or elevated despite the presence of hypercalcaemia include malignancy-associated hypercalcaemia plus concurrent primary hyperparathyroidism,[15] or patients with PTH-secreting tumours.[1] [4] 	elevated in PTH-mediated hypercalcaemia (e.g., primary hyperparathyroidism, tertiary hyperparathyroidism); suppressed in malignancy-associated hypercalcaemia unless concurrent primary hyperparathyroidism is present or ectopic PTH (rare)

Test	Result
<p>serum parathyroid hormone-related peptide</p> <ul style="list-style-type: none"> Should be ordered if initial PTH level is low or if PTH level is normal to high despite the presence of a known malignancy. Humoral hypercalcaemia of malignancy results from tumour secretion of parathyroid hormone-related peptide (PTHrP). This mechanism accounts for 80% of cases of malignancy-associated hypercalcaemia. An elevated PTHrP in combination with a suppressed PTH confirms the diagnosis. In rare cases, an elevated PTRHRP in combination with an elevated PTH suggests concurrent humoral hypercalcaemia of malignancy plus primary hyperparathyroidism.[1] [4] [5] [7] [15] 	<p>elevated in humoral hypercalcaemia of malignancy (PTHrP-mediated hypercalcaemia)</p>
<p>serum phosphorus</p> <ul style="list-style-type: none"> In humoral hypercalcaemia of malignancy, PTHrP acts at the level of the kidney to reduce calcium clearance, as well as to reduce the renal phosphorus threshold, leading to hyperphosphaturia and hypophosphataemia.[1] [4] May be included in comprehensive metabolic panel. 	<p>low in humoral hypercalcaemia of malignancy (PTHrP-mediated hypercalcaemia)</p>
<p>serum calcitriol (1,25-dihydroxyvitamin D)</p> <ul style="list-style-type: none"> Should be ordered if initial PTH level is low in the presence of lymphoma and/or granulomatous disease is suspected. Dysregulated production of calcitriol (1,25-dihydroxyvitamin D) can occur in lymphoma cells or neighboring normal cells. 	<p>elevated in calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia</p>
<p>serum 25-hydroxyvitamin D</p> <ul style="list-style-type: none"> Serum 25-hydroxyvitamin D should be checked if intravenous administration of bisphosphonate or subcutaneous denosumab is being considered, as vitamin D deficiency needs to be corrected prior to administration to avoid the risk of hypocalcaemia and, possibly, osteonecrosis of the jaw.[11] [12] [13] [14] 	<p>excludes vitamin D deficiency prior to bisphosphonate or denosumab use</p>

Other tests to consider

Test	Result
<p>skeletal survey</p> <ul style="list-style-type: none"> A skeletal survey should be done in patients in whom multiple myeloma, bone metastases, or leukaemia is suspected. 	<p>may show osteopenia, osteolytic lesions, or pathological fractures</p>
<p>chest x-ray</p> <ul style="list-style-type: none"> Hilar and/or paratracheal adenopathy with upper lobe predominant bilateral infiltrates are seen in sarcoidosis. A central mass, hilar lymphadenopathy, or pleural effusion may be seen in lung cancer. May show typical infiltrates, effusions, or cavitation in tuberculosis (TB). 	<p>may show typical findings of lung cancer, TB, or sarcoidosis</p>

Differentials

Condition	Differentiating signs / symptoms	Differentiating tests
Primary hyperparathyroidism	<ul style="list-style-type: none"> No differentiating signs or symptoms. 	<ul style="list-style-type: none"> Hypercalcaemia associated with elevated parathyroid hormone level and normal or elevated 1,25-dihydroxyvitamin D level.
Hyperthyroidism	<ul style="list-style-type: none"> Symptoms include weight loss, palpitations, anxiety, heat intolerance, hyperdefecation, tremor, diaphoresis. Signs may include a goitre or palpable thyroid nodule(s); stare; lid lag; or exophthalmos (Graves' disease). 	<ul style="list-style-type: none"> Suppressed thyroid-stimulating hormone and elevated free T4 level.
Adrenal insufficiency	<ul style="list-style-type: none"> Symptoms include fatigue, weight loss, anorexia, abdominal pain, muscle aches, lightheadedness, nausea, vomiting, and diarrhoea. Signs may include orthostatic hypotension, hyperpigmentation of skin creases and buccal mucosa. 	<ul style="list-style-type: none"> Inadequate response to adrenocorticotrophic hormone stimulation. Hyponatraemia. Hypoglycaemia.
Phaeochromocytoma	<ul style="list-style-type: none"> Symptoms include episodic headache, sweating, palpitations. Signs include persistent hypertension, tachycardia. 	<ul style="list-style-type: none"> Elevated 24-hour urinary or plasma fractionated catecholamines and metanephrines.
Sarcoidosis	<ul style="list-style-type: none"> Symptoms include cough, dyspnoea, chronic fatigue, arthralgia, wheezing, photophobia, painful red eye, blurred vision; may be asymptomatic; family history of sarcoidosis. Signs include rhonchi, lymphadenopathy, erythema nodosum, lupus pernio, conjunctival nodules, facial palsy. 	<ul style="list-style-type: none"> Chest x-ray may show hilar and/or paratracheal adenopathy with upper lobe predominant bilateral infiltrates. Serum ACE level may be elevated.
Medication-related hypercalcaemia	<ul style="list-style-type: none"> History of use of thiazide diuretics, lithium, calcium supplementation, over-the-counter antacids, or large doses of vitamin D. 	<ul style="list-style-type: none"> Serum calcium slightly elevated and returns to normal after cessation of the medications.

Condition	Differentiating signs / Differentiating tests symptoms	
Pulmonary tuberculosis	<ul style="list-style-type: none"> Cough, fever, anorexia, weight loss, and malaise. 	<ul style="list-style-type: none"> Chest x-ray: primary disease commonly presents as middle and lower lung zone infiltrates. Ipsilateral adenopathy, atelectasis from airway compression, and pleural effusion can be seen. Reactivation-type (post-primary) pulmonary tuberculosis (TB) usually involves apical and/or posterior segment of right upper lobe, apicoposterior segment of left upper lobe, or superior segment of either lower lobe, with or without cavitation. As disease progresses it spreads to other segments/lobes. Sputum smear: positive for acid-fast bacilli (AFB).
Extrapulmonary tuberculosis	<ul style="list-style-type: none"> Enlarged lymph nodes, pleuritic chest pain, skeletal pain, headache, urinary symptoms, abdominal swelling or pain. 	<ul style="list-style-type: none"> Chest x-ray: evidence of unrecognised pulmonary TB or evidence of old healed TB (e.g. upper lobe fibrosis) may be present; pleural TB will usually have a small to moderate unilateral pleural effusion. Sputum smear: positive for AFB. Analysis of lymph node aspirate; pleural, cerebrospinal or ascitic fluid; bone films; or urine may show evidence of TB infection.

Criteria

Severity

While there are no formal classification schemes for defining severity of hypercalcaemia, some consider mild hypercalcaemia to be a total adjusted serum calcium of 2.6 to 3.0 mmol/L (10.5 to 12.0 mg/dL), moderate hypercalcaemia a level of 3.0 to 3.4 mmol/L (12.0 to 13.7 mg/dL), and severe hypercalcaemia a level greater than 3.4 mmol/L (13.7 mg/dL).^{[1] [5] [6]}

Approach

Therapy directed at hypercalcaemia can temporarily restore normocalcaemia. Long-term maintenance of normocalcaemia requires eradication of the underlying malignancy. While there are no formal classification schemes for defining severity of hypercalcaemia, some consider mild hypercalcaemia to be a total adjusted serum calcium of 2.6 to 3.0 mmol/L (10.5 to 12.0 mg/dL), moderate hypercalcaemia a level of 3.0 to 3.4 mmol/L (12.0 to 13.7 mg/dL), and severe hypercalcaemia a level greater than 3.4 mmol/L (13.7 mg/dL).[1] [5] [6]

Those with mild or moderate hypercalcaemia or who are asymptomatic should repeat the serum calcium level after one week to confirm the diagnosis, be monitored, and have the underlying malignancy treated.[6]

Therapy for hypercalcaemia should be initiated for symptomatic patients and those with moderate to severe hypercalcaemia who have serum calcium concentrations >3.0 mmol/L (>12.0 mg/dL).[2] If the person has severe hypercalcaemia or severe symptoms, emergency hospital admission should be arranged.[6]

Initiation of treatment

Once the presence of hypercalcaemia has been established, initiation of treatment with intravenous normal saline, an intravenous bisphosphonate, and calcitonin can begin immediately in cases of moderate/severe or symptomatic hypercalcaemia. In cases of hypercalcaemic crisis, urgent initiation of therapy is required to achieve adequate diuresis. Once the results of additional tests are available, further targeted treatment can be started, mainly for calcitriol (1,25-dihydroxyvitamin D)-induced hypercalcaemia. For parathyroid hormone-related peptide (PTHrP)-induced hypercalcaemia, humoral hypercalcaemia, and hypercalcaemia due to metastatic skeletal involvement, no additional specific treatments are available other than treatment of the underlying malignancy.

Supportive measures

- It is important to avoid medications that can worsen hypercalcaemia (e.g., thiazide diuretics, calcitriol [(1,25-dihydroxyvitamin D)], calcium supplementation, antacids, lithium) and those that may worsen the symptoms of hypercalcaemia (e.g., sedatives, hypnotics, analgesics when possible).[1] [5]

First-line therapy

- Intravenous normal saline is one component of first-line therapy. It reverses dehydration secondary to hypercalcaemia-induced nephrogenic diabetes insipidus in addition to oral hydration, and promotes calciuresis.[1] It should be administered at 250-500 mL/hour depending on volume status, cardiac function, and kidney function.[1] [5]
- Intravenous bisphosphonates are the best-studied and most effective agents for treating malignancy-associated hypercalcaemia. Bisphosphonates effectively block osteoclastic bone resorption. Therapy should be instituted immediately upon diagnosis, because response takes 2 to 4 days.[1] Options include pamidronate disodium or zoledronic acid, which are administered as a single dose.[1] [5] [16] Based on available data, some have argued that zoledronic acid is superior to pamidronate disodium.[17] Others have suggested, however, that because the difference in control of calcaemia was small between zoledronic acid and pamidronate disodium, and the number of patients who achieved complete response by 10 days was high in both groups, either agent is acceptable.[1] Potential adverse effects associated with intravenous

bisphosphonate therapy include transient flu-like syndrome with aches/chills/fever, acute kidney injury, acute osteonecrosis of the jaw, and hypocalcaemia if high-dose bisphosphonates are given to hypercalcaemic patients with critical vitamin D deficiency.[1] [2] [4] [17] Vitamin D should be measured (and corrected if deficient) prior to intravenous bisphosphonate therapy to avoid the risk of hypocalcaemia.[2] Regular monitoring of calcium levels is also important.

Adjunctive therapies to consider

- Calcitonin interferes with osteoclastic bone resorption and renal tubular reabsorption of calcium.[1] [2] It may effect a more rapid correction of hypercalcaemia than bisphosphonates.[18] [19] The clinical utility of calcitonin is limited by its transient effect and lack of availability.[1] [5] [18] [19] Potential adverse effects include flushing and nausea.[1]
- Furosemide is reserved for managing fluid overload in conjunction with intravenous hydration. Caution should be taken to avoid overdiuresis, which depletes sodium stores relative to calcium, causing intravascular volume depletion and worsening hypercalcaemia.[5]

Recurrent or refractory hypercalcaemia

- If hypercalcaemia is improved with the initial dose, but serum calcium levels begin to increase again, the dose of bisphosphonate may be repeated after 7 days.
- In the US, denosumab is approved for the treatment of hypercalcaemia of malignancy refractory to bisphosphonate therapy. Denosumab is a monoclonal antibody directed against the receptor activator of nuclear factor-KappaB ligand (RANKL); it reduces osteoclast differentiation and bone resorption. Vitamin D should be measured (and corrected if deficient) prior to administration of denosumab to avoid the risk of hypocalcaemia. Regular monitoring of calcium levels is also important.
- In June 2018, the UK Medicines and Healthcare products Regulatory Agency (MHRA) issued a safety alert following a pooled analysis of four phase III studies of denosumab in patients with advanced malignancies involving bone.[20] New primary malignancies were reported more frequently among patients receiving denosumab than those receiving zoledronic acid (cumulative incidence of new primary malignancy at 1 year was 1.1% for denosumab and 0.6% for zoledronic acid). No treatment-related patterns for individual cancers or cancer groupings were identified. It is not known whether there is an increased risk of new primary malignancy when denosumab is prescribed for the treatment of hypercalcaemia of malignancy.

Therapy in advanced kidney disease

- Dialysis can be considered in patients who have cancers that are likely to respond to therapy, but in whom renal and cardiac function limits utilisation of intravenous hydration or accepted pharmacological therapy.[1] [21]
- Denosumab may be considered as an adjunct to dialysis.[22] However, patients with severe renal impairment (creatinine clearance <30 mL/minute) or who are receiving dialysis are at increased risk for hypocalcaemia. Regular monitoring of calcium levels is important.

Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia

Glucocorticoid therapy may be efficacious in treating calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia. No guidelines with respect to dosing and duration of therapy exist.[1]

Ectopic parathyroid hormone production

Removal of primary malignancy/ectopic parathyroid hormone-secreting source can reverse hypercalcaemia.^[23]

Treatment algorithm overview

Please note that formulations/routes and doses may differ between drug names and brands, drug formularies, or locations. Treatment recommendations are specific to patient groups: [see disclaimer](#)

Acute		(summary)
mild hypercalcaemia or asymptomatic		
	1st	treatment of underlying malignancy + supportive measures + monitoring
moderate/severe hypercalcaemia or symptomatic: without advanced kidney disease		
	1st	intravenous normal saline
	plus	intravenous bisphosphonate or denosumab
	adjunct	calcitonin (while awaiting effect of bisphosphonate)
	adjunct	furosemide
	plus	avoidance of exacerbating medications
	plus	treatment of underlying malignancy
	plus	corticosteroid
<ul style="list-style-type: none"> ■ with established calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia 		
moderate/severe hypercalcaemia or symptomatic: with advanced kidney disease		
	1st	renal dialysis ± denosumab

Treatment algorithm

Please note that formulations/routes and doses may differ between drug names and brands, drug formularies, or locations. Treatment recommendations are specific to patient groups: [see disclaimer](#)

Acute

mild hypercalcaemia or asymptomatic

1st treatment of underlying malignancy + supportive measures + monitoring

» While there are no formal classification schemes for defining severity of hypercalcaemia, some consider mild hypercalcaemia to be a total adjusted serum calcium of 2.6 to 3.0 mmol/L (10.5 to 12.0 mg/dL), moderate hypercalcaemia a level of 3.0 to 3.4 mmol/L (12.0 to 13.7 mg/dL), and severe hypercalcaemia a level greater than 3.4 mmol/L (13.7 mg/dL).[1] [5] [6]

» Those with mild or moderate hypercalcaemia or who are asymptomatic should repeat the serum calcium level after one week to confirm the diagnosis, be monitored, and have the underlying malignancy treated.[6]

» Therapy for hypercalcaemia should be initiated for symptomatic patients and those who have serum calcium concentrations >3.0 mmol/L (>12.0 mg/dL).[2]

» Long-term maintenance of normocalcaemia requires eradication of the underlying malignancy.

» It is important to avoid medications that can worsen hypercalcaemia (e.g., thiazide diuretics, calcitriol [(1,25-dihydroxyvitamin D)], calcium supplementation, antacids, lithium) and those that may worsen symptoms of hypercalcaemia (e.g., sedatives, hypnotics, analgesics, if possible).[1] [5]

» Serial monitoring of calcium is important.

moderate/severe hypercalcaemia or symptomatic: without advanced kidney disease

1st intravenous normal saline

» While there are no formal classification schemes for defining severity of hypercalcaemia, some consider mild hypercalcaemia to be a total adjusted serum calcium of 2.6 to 3.0 mmol/L (10.5 to 12.0 mg/dL), moderate hypercalcaemia a level of 3.0 to 3.4 mmol/L (12.0 to 13.7 mg/dL), and severe hypercalcaemia a level greater than 3.4 mmol/L (13.7 mg/dL).[1] [5] [6]

» Therapy for hypercalcaemia should be initiated for symptomatic patients and those who have

Acute

serum calcium concentrations >3.0 mmol/L (>12.0 mg/dL).[2]

» Once the presence of hypercalcaemia has been established, initiation of treatment with intravenous normal saline, an intravenous bisphosphonate, and calcitonin can begin immediately in cases of moderate/severe or symptomatic hypercalcaemia. In cases of hypercalcaemic crisis, urgent initiation of therapy is required to achieve adequate diuresis.

» Intravenous normal saline is one component of first-line therapy. It reverses dehydration secondary to hypercalcaemia-induced nephrogenic diabetes insipidus in addition to oral hydration, and promotes calciuresis.[1] It should be administered at 250-500 mL/hour depending on volume status, cardiac function, and kidney function.[1] [5]

plus intravenous bisphosphonate or denosumab

Treatment recommended for ALL patients in selected patient group

Primary options

» **pamidronate disodium**: 15-60 mg intravenous infusion as a single dose or given over 2-4 days (dose depends on serum calcium level, maximum 90 mg/treatment course)

OR

» **zoledronic acid**: 4 mg intravenous infusion given over at least 15 minutes; dose may be repeated after 7 days if necessary

Secondary options

» **denosumab**: consult specialist for guidance on dose

» Intravenous bisphosphonates are the best-studied and most effective agents for treating malignancy-associated hypercalcaemia. Bisphosphonates effectively block osteoclastic bone resorption. Therapy should be instituted immediately upon diagnosis, because response generally takes 2 to 4 days.[1] Options include pamidronate disodium and zoledronic acid, which are both administered as a single dose.[1] [5] [16] Based on available data, some have argued that zoledronic acid is superior to pamidronate disodium.[17] Others have suggested, however, that because the difference

Acute

in control of calcemia was small between zoledronic acid versus pamidronate disodium, and the number of patients who achieved complete response by 10 days was high in both groups, either agent is acceptable.[1] Potential adverse effects include transient flu-like syndrome with aches/chills/fever, acute kidney injury, acute osteonecrosis of the jaw, and hypocalcaemia if high-dose bisphosphonates are given to hypercalcaemic patients with critical vitamin D deficiency.[1] [2] [4] [17] Vitamin D should be measured (and corrected if deficient) prior to intravenous bisphosphonate therapy to avoid the risk of hypocalcaemia. Regular monitoring of calcium levels is also important. If hypercalcaemia is improved with the initial dose, but serum calcium levels begin to increase again, the dose may be repeated after 7 days.

» In the US, denosumab is approved for the treatment of hypercalcaemia of malignancy refractory to bisphosphonate therapy. Denosumab is a monoclonal antibody directed against the receptor activator of nuclear factor- κ B ligand (RANKL); it reduces osteoclast differentiation and bone resorption. Vitamin D should be measured (and corrected if deficient) prior to administration of denosumab to avoid the risk of hypocalcaemia. Regular monitoring of calcium levels is also important.

» In June 2018, the UK Medicines and Healthcare products Regulatory Agency (MHRA) issued a safety alert following a pooled analysis of four phase III studies of denosumab in patients with advanced malignancies involving bone.[20] New primary malignancies were reported more frequently among patients receiving denosumab than those receiving zoledronic acid (cumulative incidence of new primary malignancy at 1 year was 1.1% for denosumab and 0.6% for zoledronic acid). No treatment-related patterns for individual cancers or cancer groupings were identified. It is not known whether there is an increased risk of new primary malignancy when denosumab is prescribed for the treatment of hypercalcaemia of malignancy.

adjunct calcitonin (while awaiting effect of bisphosphonate)

Treatment recommended for SOME patients in selected patient group

Primary options

Acute

» **calcitonin-salmon**: 4-8 international units/kg intramuscularly/subcutaneously every 12 hours

» Calcitonin interferes with osteoclastic bone resorption and renal tubular reabsorption of calcium.[1] [2] It may effect a more rapid correction of hypercalcaemia than bisphosphonates.[18] [19] The clinical utility of calcitonin is limited by its transient effect and availability.[1] [5] [18] [19]

» Potential adverse effects include flushing and nausea.[1]

» Calcitonin may be used in the first 24 to 48 hours following bisphosphonate administration to transiently reverse hypercalcaemia while awaiting the therapeutic effect of bisphosphonate therapy.[24]

adjunct furosemide

Treatment recommended for SOME patients in selected patient group

Primary options

» **furosemide**: consult specialist for guidance on dose

» Furosemide is a loop diuretic reserved for managing fluid overload in conjunction with intravenous hydration. The initial dose is variable.

» Caution should be taken to avoid overdiuresis, which depletes sodium stores relative to calcium, causing intravascular volume depletion and worsening hypercalcaemia.[5]

plus avoidance of exacerbating medications

Treatment recommended for ALL patients in selected patient group

» It is important to avoid medications that can worsen hypercalcaemia (e.g., thiazide diuretics, calcitriol [(1,25-dihydroxyvitamin D)], calcium supplementation, antacids, lithium) and those that may worsen symptoms of hypercalcaemia (e.g., sedatives, hypnotics, analgesics, if possible).[1] [5]

plus treatment of underlying malignancy

Treatment recommended for ALL patients in selected patient group

» Long-term maintenance of normocalcaemia requires eradication of the underlying malignancy.

Acute

■ with established calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia

plus

» If hypercalcaemia is secondary to the rare occurrence of ectopic parathyroid hormone secretion by the underlying malignancy, removal of the primary malignancy can reverse hypercalcaemia.[23]

corticosteroid

Treatment recommended for ALL patients in selected patient group

Primary options

» prednisolone: 60 mg orally once daily for 10 days

» Once the results of additional tests are available, further targeted treatment can be started for calcitriol (1,25-dihydroxyvitamin D)-induced hypercalcaemia.

» Glucocorticoid therapy may be efficacious.

» No guidelines with respect to dosing and duration of therapy exist. However, one proposed regimen is prednisolone 60 mg/day for 10 days.[1]

moderate/severe hypercalcaemia or symptomatic: with advanced kidney disease

1st

renal dialysis ± denosumab

Primary options

» denosumab: consult specialist for guidance on dose

» Therapy for hypercalcaemia should be initiated for symptomatic patients and those who have serum calcium concentrations >3.0 mmol/L (>12.0 mg/dL).[2]

» Dialysis can be considered in patients who have cancers that are likely to respond to therapy, but in whom renal and cardiac function limits utilisation of intravenous hydration or accepted pharmacological therapy.[1] [21]

» Denosumab may be considered as an adjunct to dialysis.[22] However, patients with severe renal impairment (creatinine clearance <30 mL/minute) or who are receiving dialysis are at increased risk for hypocalcaemia. Regular monitoring of calcium levels is important.

» In June 2018, the UK Medicines and Healthcare products Regulatory Agency (MHRA) issued a safety alert following a pooled analysis of four phase III studies of denosumab in

Acute

patients with advanced malignancies involving bone.^[20] New primary malignancies were reported more frequently among patients receiving denosumab than those receiving zoledronic acid (cumulative incidence of new primary malignancy at 1 year was 1.1% for denosumab and 0.6% for zoledronic acid). No treatment-related patterns for individual cancers or cancer groupings were identified. It is not known whether there is an increased risk of new primary malignancy when denosumab is prescribed for the treatment of hypercalcaemia of malignancy.

Patient discussions

Weight-bearing ambulation and oral hydration should be encouraged. Patients should stop any calcium supplementation that they may be taking. There may need to be discussions about the continued use of medications that are associated with hypercalcaemia (e.g., lithium, thiazide diuretics, calcitriol [(1,25-dihydroxyvitamin D)], and vitamin D), and medications that may impair mental clarity (sedatives, hypnotics, analgesics).

Monitoring

Monitoring

The nadir in serum calcium levels following bisphosphonate administration typically occurs within 7 days. Responses last for 1 to 4 weeks.[17] Serial monitoring of calcium is, therefore, important. Eradication of underlying malignancy is crucial for continued maintenance of normocalcaemia.

Complications

Complications	Timeframe	Likelihood
bisphosphonate-induced flu-like syndrome	short term	low
Potential adverse effects associated with intravenous bisphosphonate therapy include transient flu-like syndrome with aches/chills/fever.[8] [9] [17]		
acute kidney injury	variable	low
Hypercalcaemia can lead to renal vasoconstriction, volume depletion, and a subsequent reversible decrease in glomerular filtration rate. Long-standing hypercalcaemia and hypercalciuria can lead to calcification, degeneration, and renal tubular atrophy/necrosis.[9] In addition, intravenous bisphosphonate therapy can be associated with acute kidney injury.[8] [9] [17]		
coma	variable	low
Hypercalcaemia is associated with neuropsychiatric symptoms, of which coma is a serious complication.		
acute pancreatitis	variable	low
Hypercalcaemia may lead to calcium deposition in the pancreatic duct, and to activation of trypsinogen in the pancreatic parenchyma, leading to pancreatitis. Incidence of acute pancreatitis associated with malignancy-associated hypercalcaemia is unknown.[10]		

Prognosis

Intravenous hydration and pharmacological therapy for malignancy-associated hypercalcaemia can provide transient restoration of normocalcaemia. Eradication of underlying malignancy is crucial for permanent reversal of hypercalcaemia.

Humoral hypercalcaemia of malignancy or local osteolytic hypercalcaemia

Intravenous hydration and calcitonin may improve hypercalcaemia within the first 24 to 48 hours of therapy.[1] [18] [19] Intravenous bisphosphonates are most efficacious and may restore normocalcaemia within 10 days of treatment in the majority of patients.[1] [17] Duration of effect is variable.[17] Sustained resolution of hypercalcaemia requires effective anti-tumour therapy.

Calcitriol (1,25-dihydroxyvitamin D)-mediated hypercalcaemia

Intravenous hydration and calcitonin may transiently improve hypercalcaemia within the first 24 to 48 hours of therapy.[1] [18] [19] Glucocorticoid therapy requires several days to improve serum calcium levels.[25] Duration of effect has not been extensively studied.

Diagnostic guidelines

United Kingdom

Emergency management of acute hypercalcaemia in adult patients (<https://www.endocrinology.org/clinical-practice/clinical-guidance/society-for-endocrinology-guidance>)

Published by: Society for Endocrinology

Last published: 2016

Clinical Knowledge Summaries: hypercalcaemia (<https://cks.nice.org.uk/hypercalcaemia>)

Published by: National Institute for Health and Care Excellence

Last published: 2019

North America

Cancer guidelines (supportive care). Oncologic emergencies: a guide for family physicians (<https://www.albertahealthservices.ca/info/cancerguidelines.aspx>)

Published by: Alberta Provincial Tumour Council

Last published: 2014

Treatment guidelines

United Kingdom

Emergency management of acute hypercalcaemia in adult patients (<https://www.endocrinology.org/clinical-practice/clinical-guidance/society-for-endocrinology-guidance>)

Published by: Society for Endocrinology

Last published: 2016

Clinical Knowledge Summaries: hypercalcaemia (<https://cks.nice.org.uk/hypercalcaemia>)

Published by: National Institute for Health and Care Excellence

Last published: 2019

International

Hypercalcaemia of malignancy (HCM) (<https://www.eviq.org.au/clinical-resources/oncological-emergencies/486-hypercalcaemia-of-malignancy-hcm#management>)

Published by: Cancer Institute NSW, New South Wales Government, Australia

Last published: 2019

North America

Cancer guidelines (supportive care). Oncologic emergencies: a guide for family physicians (<https://www.albertahealthservices.ca/info/cancerguidelines.aspx>)

Published by: Alberta Provincial Tumour Council

Last published: 2014

Key articles

- Stewart AF. Clinical practice. Hypercalcemia associated with cancer. *N Engl J Med*. 2005 Jan 27;352(4):373-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com)
- Horwitz MJ. Chapter 84: Non-parathyroid hypercalcemia. In: Bilezikian JP, ed. *Primer on the metabolic bone diseases and disorders of mineral metabolism*. 9th ed. Washington, DC: American Society of Bone and Mineral Research; 2018:639-45.
- Alberta Provincial Tumour Council. Oncologic emergencies: a guide for family physicians. Sep 2014 [internet publication]. [Full text \(https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-oncologic-emergencies.pdf\)](https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-oncologic-emergencies.pdf)
- Stewart AF. Clinical practice. Hypercalcemia associated with cancer. *N Engl J Med*. 2005 Jan 27;352(4):373-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com)
- Horwitz MJ, Hodak SP, Stewart AF. Non-parathyroid hypercalcemia. In: Rosen CJ, ed. *Primer on the metabolic bone diseases and disorders of mineral metabolism*. 8th ed. Washington, DC: American Society of Bone and Mineral Research; 2013:562-71.
- Alberta Provincial Tumour Council. Oncologic emergencies: a guide for family physicians. Sep 2014 [internet publication]. [Full text \(https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-oncologic-emergencies.pdf\)](https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-oncologic-emergencies.pdf)
- Body JJ, Bartl R, Burckhardt P, et al. Current use of bisphosphonates in oncology. International Bone and Cancer Study Group. *J Clin Oncol*. 1998 Dec;16(12):3890-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9850035?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9850035?tool=bestpractice.bmj.com)
- Major P, Lortholary A, Hon J, et al. Zoledronic acid is superior to pamidronate in the treatment of hypercalcemia of malignancy: a pooled analysis of two randomized, controlled clinical trials. *J Clin Oncol*. 2001 Jan 15;19(2):558-67. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11208851?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11208851?tool=bestpractice.bmj.com)

References

1. Stewart AF. Clinical practice. Hypercalcemia associated with cancer. *N Engl J Med*. 2005 Jan 27;352(4):373-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com)
2. Cancer Institute NSW. Hypercalcaemia of malignancy (HCM). Jul 2019 [internet publication]. [Full text \(https://www.eviq.org.au/clinical-resources/oncological-emergencies/486-hypercalcaemia-of-malignancy-hcm\)](https://www.eviq.org.au/clinical-resources/oncological-emergencies/486-hypercalcaemia-of-malignancy-hcm)
3. Lindner G, Felber R, Schwarz C, et al. Hypercalcemia in the ED: prevalence, etiology, and outcome. *Am J Emerg Med*. 2013 Apr;31(4):657-60. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/23246111?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/23246111?tool=bestpractice.bmj.com)

4. Horwitz MJ. Chapter 84: Non-parathyroid hypercalcemia. In: Bilezikian JP, ed. Primer on the metabolic bone diseases and disorders of mineral metabolism. 9th ed. Washington, DC: American Society of Bone and Mineral Research; 2018:639-45.
5. Alberta Provincial Tumour Council. Oncologic emergencies: a guide for family physicians. Sep 2014 [internet publication]. Full text (<https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-oncologic-emergencies.pdf>)
6. National Institute for Health and Care Excellence. Clinical Knowledge Summaries: hypercalcaemia. Aug 2019 [internet publication]. Full text (<https://cks.nice.org.uk/hypercalcaemia>)
7. Strewler GJ, Budayr AA, Clark OH, et al. Production of parathyroid hormone by a malignant nonparathyroid tumor in a hypercalcemic patient. J Clin Endocrinol Metab. 1993 May;76(5):1373-5. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/7684395?tool=bestpractice.bmj.com>)
8. Stewart AF. Clinical practice. Hypercalcemia associated with cancer. N Engl J Med. 2005 Jan 27;352(4):373-9. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/15673803?tool=bestpractice.bmj.com>)
9. Horwitz MJ, Hodak SP, Stewart AF. Non-parathyroid hypercalcemia. In: Rosen CJ, ed. Primer on the metabolic bone diseases and disorders of mineral metabolism. 8th ed. Washington, DC: American Society of Bone and Mineral Research; 2013:562-71.
10. Alberta Provincial Tumour Council. Oncologic emergencies: a guide for family physicians. Sep 2014 [internet publication]. Full text (<https://www.albertahealthservices.ca/assets/info/hp/cancer/if-hp-cancer-guide-oncologic-emergencies.pdf>)
11. Lorenzo-Pouso AI, Pérez-Sayáns M, García A, et al. Vitamin D supplementation: hypothetical effect on medication-related osteonecrosis of the jaw. Med Hypotheses. 2018 Jul;116:79-83. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/29857915?tool=bestpractice.bmj.com>)
12. Bedogni A, Bettini G, Bedogni G, et al. Is vitamin D deficiency a risk factor for osteonecrosis of the jaw in patients with cancer? A matched case-control study. J Craniomaxillofac Surg. 2019 Aug;47(8):1203-8. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/30929994?tool=bestpractice.bmj.com>)
13. Daga N, Joseph F. Denosumab-induced severe hypocalcaemia in a patient with vitamin D deficiency. BMJ Case Rep. 2020 Aug 26;13(8):e234508. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/32847872?tool=bestpractice.bmj.com>)
14. Râkel A, Boucher A, Ste-Marie LG. Role of zoledronic acid in the prevention and treatment of osteoporosis. Clin Interv Aging. 2011;6:89-99. Full text (<https://www.doi.org/10.2147/CIA.S7282>) Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/21594000?tool=bestpractice.bmj.com>)
15. Milanesi A, Yu R, Geller SA, et al. Concurrent primary hyperparathyroidism and humoral hypercalcemia of malignancy in a patient with multiple endocrine neoplasia type 1. Pancreas. 2011 May;40(4):634-7. Abstract (<http://www.ncbi.nlm.nih.gov/pubmed/21483254?tool=bestpractice.bmj.com>)

16. Body JJ, Bartl R, Burckhardt P, et al. Current use of bisphosphonates in oncology. International Bone and Cancer Study Group. *J Clin Oncol*. 1998 Dec;16(12):3890-9. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/9850035?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/9850035?tool=bestpractice.bmj.com)
17. Major P, Lortholary A, Hon J, et al. Zoledronic acid is superior to pamidronate in the treatment of hypercalcemia of malignancy: a pooled analysis of two randomized, controlled clinical trials. *J Clin Oncol*. 2001 Jan 15;19(2):558-67. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/11208851?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/11208851?tool=bestpractice.bmj.com)
18. Ljunghall S. Use of clodronate and calcitonin in hypercalcemia due to malignancy. *Recent Results Cancer Res*. 1989;116:40-5. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/2527399?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/2527399?tool=bestpractice.bmj.com)
19. Chevallier B, Peyron R, Basuyau JP, et al. Human calcitonin in neoplastic hypercalcemia. Results of a prospective randomized trial [in French]. *Presse Med*. 1988 Dec 17;17(45):2375-7. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/2974978?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/2974978?tool=bestpractice.bmj.com)
20. Medicines and Healthcare products Regulatory Agency. Denosumab (Xgeva) for advanced malignancies involving bone: study data show new primary malignancies reported more frequently compared to zoledronate. Jun 2018 [internet publication]. [Full text \(https://www.gov.uk/drug-safety-update/denosumab-xgeva-for-advanced-malignancies-involving-bone-study-data-show-new-primary-malignancies-reported-more-frequently-compared-to-zoledronate\)](https://www.gov.uk/drug-safety-update/denosumab-xgeva-for-advanced-malignancies-involving-bone-study-data-show-new-primary-malignancies-reported-more-frequently-compared-to-zoledronate)
21. Koo WS, Jeon DS, Ahn SJ, et al. Calcium-free hemodialysis for the management of hypercalcemia. *Nephron*. 1996;72(3):424-8. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/8852491?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/8852491?tool=bestpractice.bmj.com)
22. Bech A, de Boer H. Denosumab for tumor-induced hypercalcemia complicated by renal failure. *Ann Intern Med*. 2012 Jun 19;156(12):906-7. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/22711097?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/22711097?tool=bestpractice.bmj.com)
23. Nussbaum SR, Gaz RD, Arnold A. Hypercalcemia and ectopic secretion of parathyroid hormone by an ovarian carcinoma with rearrangement of the gene for parathyroid hormone. *N Engl J Med*. 1990 Nov 8;323(19):1324-8. [Full text \(https://www.nejm.org/doi/full/10.1056/NEJM199011083231907\)](https://www.nejm.org/doi/full/10.1056/NEJM199011083231907) [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/2215618?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/2215618?tool=bestpractice.bmj.com)
24. Wisneski LA. Salmon calcitonin in the acute management of hypercalcemia. *Calcif Tissue Int*. 1990;46 Suppl:S26-30. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/2137363?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/2137363?tool=bestpractice.bmj.com)
25. Binstock ML, Mundy GR. Effect of calcitonin and glucocorticoids in combination on the hypercalcemia of malignancy. *Ann Intern Med*. 1980 Aug;93(2):269-72. [Abstract \(http://www.ncbi.nlm.nih.gov/pubmed/7406378?tool=bestpractice.bmj.com\)](http://www.ncbi.nlm.nih.gov/pubmed/7406378?tool=bestpractice.bmj.com)

Images




Figure 1: CT chest showing compression fracture of multiple vertebral bodies in a child presenting with acute lymphoblastic leukaemia. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level

Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758




Figure 2: Whole body planar images suggestive of skeletal infiltration in a child with acute lymphoblastic leukaemia showing areas of abnormal increased uptake. Biochemistry showed hypercalcaemia with a suppressed parathyroid hormone level

Sukumar SP, Balachandran K, Sahoo JP, et al. Acute lymphocytic leukaemia presenting as a metabolic bone disease. BMJ Case Reports 2013; doi:10.1136/bcr-2013-008758

Disclaimer

BMJ Best Practice is intended for licensed medical professionals. BMJ Publishing Group Ltd (BMJ) does not advocate or endorse the use of any drug or therapy contained within this publication nor does it diagnose patients. As a medical professional you retain full responsibility for the care and treatment of your patients and you should use your own clinical judgement and expertise when using this product.

This content is not intended to cover all possible diagnosis methods, treatments, follow up, drugs and any contraindications or side effects. In addition, since such standards and practices in medicine change as new data become available, you should consult a variety of sources. We strongly recommend that you independently verify specified diagnosis, treatments and follow-up and ensure it is appropriate for your patient within your region. In addition, with respect to prescription medication, you are advised to check the product information sheet accompanying each drug to verify conditions of use and identify any changes in dosage schedule or contraindications, particularly if the drug to be administered is new, infrequently used, or has a narrow therapeutic range. You must always check that drugs referenced are licensed for the specified use and at the specified doses in your region.

Information included in BMJ Best Practice is provided on an “as is” basis without any representations, conditions or warranties that it is accurate and up to date. BMJ and its licensors and licensees assume no responsibility for any aspect of treatment administered to any patients with the aid of this information. To the fullest extent permitted by law, BMJ and its licensors and licensees shall not incur any liability, including without limitation, liability for damages, arising from the content. All conditions, warranties and other terms which might otherwise be implied by the law including, without limitation, the warranties of satisfactory quality, fitness for a particular purpose, use of reasonable care and skill and non-infringement of proprietary rights are excluded.

Where BMJ Best Practice has been translated into a language other than English, BMJ does not warrant the accuracy and reliability of the translations or the content provided by third parties (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages). BMJ is not responsible for any errors and omissions arising from translation and adaptation or otherwise. Where BMJ Best Practice lists drug names, it does so by recommended International Nonproprietary Names (rINNs) only. It is possible that certain drug formularies might refer to the same drugs using different names.

Please note that recommended formulations and doses may differ between drug databases drug names and brands, drug formularies, or locations. A local drug formulary should always be consulted for full prescribing information.

Treatment recommendations in BMJ Best Practice are specific to patient groups. Care is advised when selecting the integrated drug formulary as some treatment recommendations are for adults only, and external links to a paediatric formulary do not necessarily advocate use in children (and vice-versa). Always check that you have selected the correct drug formulary for your patient.

Where your version of BMJ Best Practice does not integrate with a local drug formulary, you should consult a local pharmaceutical database for comprehensive drug information including contraindications, drug interactions, and alternative dosing before prescribing.

Interpretation of numbers

Regardless of the language in which the content is displayed, numerals are displayed according to the original English-language numerical separator standard. For example 4 digit numbers shall not include a comma nor a decimal point; numbers of 5 or more digits shall include commas; and numbers stated to be less than 1 shall be depicted using decimal points. See Figure 1 below for an explanatory table.

BMJ accepts no responsibility for misinterpretation of numbers which comply with this stated numerical separator standard.

This approach is in line with the guidance of the [International Bureau of Weights and Measures Service](#).

Figure 1 – BMJ Best Practice Numeral Style

5-digit numerals: 10,000

4-digit numerals: 1000

numerals < 1: 0.25

Our full website and application terms and conditions can be found here: [Website Terms and Conditions](#).

Contact us

+ 44 (0) 207 111 1105

support@bmj.com

BMJ

BMA House

Tavistock Square

London

WC1H 9JR

UK

BMJ Best Practice

Contributors:

// Authors:

Albert Shieh, MD

Assistant Professor of Medicine

Division of Geriatric Medicine, Department of Medicine, University of California, Los Angeles, Los Angeles, CA

DISCLOSURES: AS declares that he has no competing interests.

Dorothy Martinez, MD

Clinical Professor of Medicine

Division of Endocrinology, Department of Medicine, University of California, Los Angeles, Los Angeles, CA

DISCLOSURES: DM declares that she has no competing interests.

// Peer Reviewers:

Ken C. Chiu, MD, FACE

Professor

Endocrinology Fellowship Training Program, Department of Clinical Diabetes, Endocrinology, and Metabolism, City of Hope National Medical Center, Duarte, CA

DISCLOSURES: KCC declares that he has no competing interests.

Richard Quinton, MD

Consultant and Senior Lecturer

Endocrine Unit, Newcastle University and Royal Victoria Infirmary, Newcastle-upon-Tyne, UK

DISCLOSURES: RQ has been reimbursed by Amgen, the manufacturer of Cinacalcet, for attending a conference and for giving an educational seminar (total under £1,200 over past 3 years).